

Title: Chronic Granulomatous Disease *GeneReview* – Research Testing

Authors: Leiding JW, Holland SM

Updated: February 2016

Research Testing

Determination of residual superoxide function is important in management: patients with little to no superoxide production are at the greatest risk for mortality [Kuhns et al 2010] and, thus, are the most likely candidates for HSCT. The following three tests are performed in research laboratories only:

- Cytochrome *c* reduction assay quantitates indirectly the actual amount of superoxide produced by measuring spectrophotometrically the inhibitable reduction of ferricytochrome *c* by superoxide dismutase to ferrocycytochrome *c* [Elloumi & Holland 2014]. Results of this test correlate well with results of the DHR test.
- Chemiluminescence. Because superoxide can cause a variety of chemical agents to luminesce, measurement of luminescence (typically using dichlorofluorescein [DCF]) can quantitate the amount of superoxide produced [Elloumi & Holland 2014]. While this assay can rapidly detect superoxide activity and identify hypomorphic forms of CGD, it lacks cellular resolution and thus cannot identify female carriers of X-linked CGD.
- Neutrophil superoxide production of reactive oxygen intermediates (ROI). The quantitation of superoxide produced can be obtained directly from the cytochrome *c* reduction assay (a research laboratory test) or indirectly from the DHR test (a routine clinical test). In general, a DHR test value in the lower range (i.e., <225 arbitrary units) correlates with poor superoxide production, which can be predicted from the specific NADPH oxidase pathogenic variant (see [Genotype-Phenotype Correlations](#)).

Immunoblot test for the NADPH complex proteins. Failure to detect the following cytoplasmic subunits of the phagocyte NADPH oxidase (phox) proteins suggests autosomal recessive inheritance: p47^{phox} (encoded by *NCF1*), p67^{phox} (*NCF2*), or p40^{phox} (*NCF4*) ([Table 1](#) and [Table A](#)). Immunoblotting is currently performed only in research laboratories.

Note: This technique cannot distinguish between pathogenic variants in [CYBB](#) (encoding gp91^{phox}) and [CYBA](#) (encoding p22^{phox}). Because the protein products of these two genes stabilize each other within the phagocyte membrane, absence of one protein results in the absence of the other [Segal et al 2000] (see [Molecular Genetics](#)). Pathogenic variants in *CYBB* or *CYBA* that cause a failure to bind heme (leading to a loss of the cytochrome b558) have been referred to as cytochrome negative. In contrast, pathogenic variants in *NCF1*, *NCF2*, and *NCF4* leave cytochrome b558 intact and have been referred to as cytochrome positive. Because pathogenic missense variants in either *CYBB* or *CYBA* can also support cytochrome b558 persistence without function, the terminology ‘cytochrome negative’ and ‘cytochrome positive’ is not preferred.

References

Elloumi HZ, Holland SM. Diagnostic assays for chronic granulomatous disease and other neutrophil disorders. *Methods Mol Biol.* 2014;1124:517-35.

Kuhns DB, Alvord WG, Heller T, Feld JJ, Pike KM, Marciano BE, Uzel G, Deravin SS, Priel DA, Soule BP, Zarembek KA, Malech HL, Holland SM, Gallin JI. Residual NADPH oxidase and survival in chronic granulomatous disease. *N Engl J Med.* 2010;363:2600-10.

Segal BH, Leto TL, Gallin JI, Malech HL, Holland SM. Genetic, biochemical, and clinical features of chronic granulomatous disease. *Medicine (Baltimore).* 2000;79:170-200.